

Gray whale's body condition in Mexico 2020




Floryser Ronzón, Sergio Martínez, Lorena Vilorio, Steven Swartz, Jorge Urbán

Survey area

LSI 2008-2011
2018-2020








BM 2018-2020



Body condition in live gray whales

GOOD	322	321	32X	312	31X	3X2	3X1	3XX		
FAIR	311	222	221	22X	212	21X	2X2	2X1	2XX	
POOR	211	122	121	12X	112	111	11X	1X2	1X1	1XX
UNKNOWN	X22	X21	X2X	X12	X11	X1X	XX2	XX1	XXX	

X=No photo

	Head	Scapula	Flank
3			
2			
1			

Body condition in live gray whales




GOOD

FAIR

POOR


Body condition in live gray whales in LSI (Single whales)




n=	89	236	433	347	207	529	499
Poor Condition	6.7%	7.6%	6.2%	4.9%	8.2%	23.6%	30%




Body condition in live gray whales in BM (Single whales)



Female /calf with photo-ID in LSI and BM by year




Body condition in live gray whales in LSI (Female with calf)



	2008	2009	2010	2011	2018	2019	2020
n=	79	70	31	176	80	40	54
Poor Condition	0%	0%	0%	2.3%	2.5%	0%	6%

Body condition in live gray whales in BM (Female with calf)



n=
Poor
Condition

27	13	24
4%	8%	0%